



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

## HOW TO DETERMINE THE PRESENCE OF INTESTINAL PARASITES

Darling (Société de Pathologic Exotique, May 10, 1911) describes two simple methods for determining the presence of intestinal parasites, especially the oxyuris, *Ankylostoma duodenale* and the *Strongylus stercorale*.

The first method consists in mixing a small portion of the feces with sterile water in a Petri dish and incubating at 37 degrees for twenty-four hours.

The second method is based upon the separation of the ova by means of centrifuging. The specimen is shaken vigorously in a test tube with sterilized distilled water, then centrifuged after calcium chloride has been added. The supernatant fluid is then examined for the ova.—(V. A. L., in *La Tribune Med.*, Mch., 1912.)

## EXAMINATION OF SPUTUM FOR TUBERCLE BACILLI

Many methods have been tested by which sputum may be liquified and the bacilli precipitated, and the use of the centrifuge and filter dispensed with. The following are presented by Nemmsner and Martos-Lissowska (*Deutsch Med. Woch.*, Sept. 14, 1911; p. 1697);

(1) *Alkaline trypsin digestion.*

To 0.1 cc. of trypsin and 5 cc. of a 0.4% sodium hydrate solution are added 5 cc. of sputum. Shake the mixture with the addition of a few drops of  $\text{CHCl}_3$  and incubate at 37° C for 24 hours. Then there will be found beneath a clear supernatant liquid, a *compact* deposit, which can be readily removed and stained.\*

(2) *Acid trypsin digestion.*

As above; except that 0.4% HCl is substituted for the soda solution, and no  $\text{CHCl}_3$  need be added.

(3) *Oxidation.*

To 5 cc. of sputum 5-10 drops of perchloric acid— $\text{HClO}_4$ —and 5 cc. of water are added. This is incubated, and treated as before. Instead of the perchloric acid, 0.5 gm. of potassium chlorate and 5 cc. of 0.4% HCl may be substituted.

\*Much's method of sputum staining is very valuable for all suspected cases of phthisis in which no tubercle bacilli are found by the usual methods of examination. This consists of treatment with anilin gentian violet for 48 hours and then with Gram's solution. By this means non-acid fast forms of tubercle bacilli, which fail to retain the dye under Ziehl's process, may be seen. See Roepke, *Deutsch Med. Woch.*, Oct., 1911, p. 1937, or J. R. A. and Med. Corps, Mch., 1912, p. 357.